

REFERENCES

- Affonso, D., Bosque, E., Wahlberg, V., & Brady, J.P. (1993). Reconciliation and healing of mothers through skin-to-skin contact provided in an American tertiary level intensive care unit. *Neonatal Network*, 12(3), 25-32.
- Aita, M., & Snider, L. (2003). The art of developmental care in the NICU: a conceptual analysis. *Journal of Advanced Nursing*, 41(3), 223-232.
- Als, H. (1982). Towards a synactive theory of development: promise for the assessment and support of infant individuality. *Infant Mental Health Journal*, 3(4), 229-243.
- Als, H. (1984). Newborn Behavioral Assessment. Progress in Pediatric Psychology. New York: Grune & Stratton, 1-46.
- Als, H. (1986). A synactive model of neonatal behavioral organization: framework for the assessment of neurobehavioral development in the premature infant and for support of infants and parents in the neonatal intensive care environment. *Physical and Occupational Therapy in Pediatrics*, 6(3/4), 3-55.
- Als, H., Lawhon, G., Duffy, F.H., McAnulty, G.B., Gibes-Grossman, R., & Blickman, J.G. (1994). Individualized developmental care for the very low birth weight preterm infant: medical and neurofunctional effects. *Journal of American Medical Association*, 272, 853-858.
- Als, H., & Gilkerson, L. (1995). Developmentally supportive care in the neonatal intensive care unit. *Zero to Three*, 15(6), 1-10.
- Als, H., & Gilkerson, L. (1997). The role of relationship based developmentally supportive newborn intensive care in strengthening outcome of preterm infants. *Seminars in Perinatology*, 21(3), 178-179.
- Als, H. (1997). Manual for the Naturalistic Observation of Newborn Behavior (NIDCAP®). Boston: National NIDCAP® Center. Children's Hospital.
- Als, H. (1998). Developmental care in the newborn intensive care unit. *Current Opinion in Pediatrics*, 10, 138-142.
- Als, H., & Gilkerson, L., Duffy, F.H., McAnulty, G.B., et al. (2003). A three-center, randomized, controlled trial of individualized developmental care for very low birth weight preterm infants: medical, neurodevelopmental, parenting and caregiving effects. *Journal of Developmental and Behavioral Pediatrics*, 24 (6), 399-408.
- Als, H., Duffy, F.H., McAnulty, G.B., Rivkin, M.J., Vajapeyam, S., Mulkern, R.V., Warfield, S.K., Hupp, P.S., Butler, S., Conneman, N., Fischer, C., & Eichenwald, E.C. (2004). Early experience alters brain function and structure. *Pediatrics*, 113 (4), 846-857.
- Anand, K.J.S., & Scalzo, F.M. (2000). Can adverse neonatal experiences alter brain development and subsequent behavior? *Biology of the Neonate*, 77, 69-82.
- Anderson, G.C., Lane, A.E., & Chang, H.P. (1995). Axillary temperature in the transitional newborn infants before and after tub bath. *Applied Nursing Research*, 8(3), 123-128.
- Buehler, D.M., Als, H., Duffy, F.H., McAnulty, G.B., & Liederman, J. (1995). Effectiveness of individualized developmental care for low-risk preterm infants: behavioral and electrophysiologic evidence. *Pediatrics*, 96(5), 923-932.
- Bremner, P., Byers, J., & Kiehl, E. (2003). Noise and the premature infant: physiological effects and practice implications. *Journal of Obstetric, Gynecologic and Neonatal Nursing*, 32(4), 447-454.
- Bullough, J., & Rea, M. (1996). Lighting for neonatal intensive care units: Some Critical information for design. *Lighting Research Technology*, 28, 189-198.
- Blackburn, S. (1998). Environmental impact of the NICU on developmental outcomes. *Journal of Pediatric Nursing*, 13(5), 279-289.
- Blackburn, S., & Peterson, D. (1991). Effects of cycled lighting on activity state and cardiorespiratory function in preterm neonates. *Journal of Perinatal Neonatal Nursing*, 4(4), 47-54.
- Brandon, D.H., Holditch-Davis, D., & Belyea, M. (2002). Preterm infants born at less than 31 weeks' gestation have improved growth in the cycled light compared with continuous near darkness. *Journal of Pediatrics*, 140, 192-199.
- Catlett, A.T., & Holditch-Davis, D. (1990). Environmental stimulation of the acutely ill premature infant: physiological effects and nursing implications. *Neonatal Network*, 8(6), 19-26.
- Cetta, F., Lambert, G.H., & Ross, P. (1991). Newborn chemical exposure from over-the-counter skin care products. *Clinical Pediatrics (Phila)*, 30(5), 286-289.
- Chang, Y., Anderson, G.C., & Lin, C. (2002). Decreased activity and oxygen desaturation in prone ventilated preterm infants during the first postnatal week. *Heart & Lung*, 31(1), 34-42.
- Chang, Y., Anderson, G., & Lin, C. (2002). Effects of prone and supine positions on sleep states and stress responses in mechanically ventilated preterm infants during the first postnatal week. *Journal of Advanced Nursing*, 40(2), 161-169.
- Chwo, M., Anderson, G., & Good, M. (2002). A randomized controlled trial of early kangaroo care for preterm infants: effects on temperature, weight, and acuity. *Journal of Nursing Research*, 10(2), 129-142.
- Cole, J.J., Begish-Duddy, A., Judas, M.L., & Jorgensen, K. M. (1990). Changing the NICU environment: the Boston city hospital model. *Neonatal Network*, 9(2), 15-23.
- Cole, P., & Gavey, J. (2001). Are they lying comfortably? A direct observational study examining current practices of infant positioning in the neonatal unit. *Journal of Neonatal Nursing*, 7(1), 25-28.
- Cowan, M.E., & Frost, M.R. (1986). A comparison between a detergent baby bath additive and baby soap on the skin flora of neonates. *Journal of Hospital Infection*, 7(1), 91-95.
- Darmstadt, G.L., & Dinulos, J.G. (2000). Neonatal skin care. *Pediatric Clinics of North America*, 47(4), 757-782. Review.
- Ewer, A., James, M., & Tobin, J. (1999). Prone and left lateral positioning reduce gastro-oesophageal reflux in preterm infants. *Archives of Disease in Childhood: Fetal & Neonatal Ed.*, 81(3), F201-5.
- Eichler, F., Ipsiroglu, O., Arif, T., et al. (2001). Position dependent changes of cerebral blood flow velocities in premature infants. *European Journal of Pediatrics*, 160(10), 633-639.
- Fern, D., Graves, C., & L'Huillier, M. (2002). Swaddled bathing in the newborn intensive care unit. *Newborn and Infant Nursing Reviews*, 44(4), 1093-1118.
- Fleisher, B.F., VandenBerg, K.A., Constantinou, J., et al. (1995). Individualized developmental care for very low birth weight premature infants. *Clinical Pediatrics (Phila)*, 34, 522-529.
- Fox, M., & Molesky, M. (1990). The effects of prone and supine positioning on arterial oxygen pressure. *Neonatal Network*, 8(4), 25-29.
- Frank, L.S., Quinn, D., & Zahr, L. (2000). Effect of less frequent bathing of preterm infants on skin flora and pathogen colonization. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 29(6), 584-589.
- Gale, G., & VandenBerg, K.A. (1998). Kangaroo Care. *Neonatal Network*, 17(5), 69-71.
- Gfatter, R., Hackl, P., & Braun, F. (1997). Effects of soap and detergents on skin surface pH, stratum corneum hydration and fat content in infants. *Dermatology*, 195(3), 258-262.
- Glass, P. (1993). Development of visual function on preterm infants: implications of early intervention. *Infants and Young Children*, 6(1), 11-20.
- Grenier, I., Bigsby, R., Vergara, E., et al. (2003). Comparison of motor, self-regulatory, and stress behaviors of preterm infants across body positions. *American Journal of Occupational Therapy*, 57(3), 289-297.
- Gressens, P., Rogido, M., Painsaveine, B., & Sola, A. (2002). The impact of neonatal intensive care practices on the developing brain. *Journal of Pediatrics*, 140(6), 646-653.
- Grunau, R., Linhares, M., Holsti, M., et al. (2004). Does prone or supine position influence pain responses in preterm infants at 32 weeks gestational age? *Clinical Journal of Pain*, 20(2), 76-82.
- Hadley, L.B., West, W., Turner, A., & Santangelo, S. (1999). *Developmental and behavioral characteristics of preterm infants*. Santa Rosa, CA: NICU INK.
- Hanson, J.L., Jepson, E.S., Johnson, B.H., & Thomas, J. (1997). *Family-centered care: Changing practice, changing attitudes*. Bethesda, MD: Institute for Family-Centered Care.
- Heller, A. (1991). A comparison of the effects of containment and stroking of preterm infants at varying levels of maturity. Chicago: Loyola University of Chicago, Dissertation.
- Holditch-Davis, D., & Miles, M. (2000). Mother's stories about their experiences in the neonatal intensive care unit. *Neonatal Network*, 19(3), 3-21.
- Horton, J., Waldertstrom, U., & Bowman, E. (1998). Touch of low birth weight babies in neonatal intensive care: Observations over a 24 hour period. *Journal of Neonatal Nursing*, 4, 24-28.
- Jackson, K., Ternestedt, B., & Schollin, J. (2003). From alienation to familiarity: experiences of mothers and fathers of preterm infants. *Journal of Advanced Nursing*, 43(2), 120-129.
- Jacobs, S., Sokol, J., & Ohlsson, A. (2002). The newborn individualized developmental care and assessment program is not supported by meta-analyses of the data. *Journal of Pediatrics*, 140, 699-706.
- Johnson, A. (2003). Adapting the neonatal intensive care environment to decrease noise. *Journal of Perinatal and Neonatal Nursing*, 17(4), 280-290.
- Johnson, B.H. (1995). Newborn intensive care units pioneer family-centered change in hospitals across the country. *Zero to Three*, 15, 11-17.
- Johnston, C., Stevens, B., Pinelli, J., et al. (2003). Kangaroo care is effective in diminishing pain response in preterm neonates. *Archives of Pediatric and Adolescent Medicine*, 157(11), 1084-1088.
- Jorgensen, K.M. (1995). Infant Skin pH: How soaps affect the delicate balance. *NeoNotes*.
- Kenner, C., & McGrath, J. (2004). *Developmental care of newborns and infants: A guide for health professionals*. St. Louis, MO: National Association of Neonatal Nurses.
- Kleberg, A., Westrup, B., Stjernqvist, K., & Lagercrantz, H. (2002). Indications of improved cognitive development at one year of age among infants born very prematurely who received care based on the Newborn Individualized Developmental Care and Assessment Program (NIDCAP). *Early Human Development*, 68, 83-91.
- Law-Morstatt, L., Judd, D.M., Snyder, P., Baier, R.J., & Dhanireddy, R. (2003). Pacing as a treatment technique for transitional sucking patterns. *Journal of Perinatology*, 23(6), 483-488.
- Lee, H. K. (2002). Effects of sponge bathing on vagal tone and behavioral responses in premature infants. *Journal of Clinical Nursing*, 11(4), 510-519.
- Legault, M., & Goulet, C. (1995). Comparison of kangaroo care and traditional methods of removing preterm infants from incubators. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 24(6), 501-506.
- Lotas, M.J. (1992). Effects of light and sound in the neonatal intensive care environment on the low birth weight infant. *NAACOG's Clinical Issues*, 3(1), 34-44.
- Ludington-Hoe, S., & Swinsh, J. (1996). Developmental aspects of kangaroo care. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 25(8), 691-703.
- Lund, C., Kuller, J., Lane, A., Lott, J.W., & Raines, D.A. (1999). Neonatal skin care: The scientific basis for practice. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 28, 241-254.
- Lund, C., Kuller, J., Lane, A., Lott, J.W., Osborne, J.W., & Raines, D.A. (2001). Neonatal skin care: Clinical outcomes of the AWHONN/NANN evidence-based clinical practice guidelines. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 30, 41-51.
- Maynard, V., Bignall, S., & Kitchen, S. (2000). Effect of positioning on respiratory synchrony in non-ventilated preterm infants. *Physiotherapy Research International*, 5(2), 96-110.
- McCain, G.C. (2003). An evidence-based guideline or introducing oral feeding to healthy preterm infants. *Neonatal Network*, 22(5).
- McCain, G.C., & Gartside, P.S. (2002). Behavioral responses of preterm infants to a standard-care and semi-demand feeding protocol. *Newborn and Infant Nursing Reviews*, 2, 187-193.
- McGrath, J.M., & Medoff-Cooper, B. (2002). Alertness and feeding competence in extremely early born preterm infants. *Newborn and Infant Nursing Reviews*, 2(3), 174-186.
- Miller, C.L., White, R., Whitman, T. L., O'Callaghan, M.F., & Maxwell, S.E. (1995). The effects of cycled versus non-cycled lighting on growth and development in preterm infants. *Infant Behavior and Development*, 18, 87-95.
- Mirmiran, M., & Ariago, R.L. (2000). Influence of light in the NICU on the developmental of circadian rhythms in preterm infants. *Seminars in Perinatology*, 24, 247-257.
- Moore, K., Coker, K., DuBuisson, A., Swett, B., & Edwards, W. (2003). Implementing potentially better practices for improving family-centered care in neonatal intensive care units: successes and challenges. *Pediatrics*, 111(4), 450-460.
- Morelli, J.G., & Weston, W.L. (1987). Soaps and shampoos in pediatric practice. *Pediatrics*, 80(5), 634-637.
- Munson, K.A., Bare, D.E., Hoath, S.B., & Visscher, M.O. (1999). A survey of skin care practices for premature low birth weight infants. *Neonatal Network*, 18(3), 25-31.
- Peters, K.L. (1998). Bathing premature infants: physiological and behavioral consequences. *American Journal of Critical Care*, 7(2), 90-100.
- Phelps, D. L., & Watts, J.L. (2000). Early light reduction for prevention of retinopathy of prematurity in very low birth weight infants. *Cochrane Database System Review*, (2), CD000122.
- Premji, S.S., Chessell, L., Paes, B., Pinelli, J., & Jacobson, K. (2002). A matched cohort of feeding practice guidelines for infants weighing less than 1500g. *Advances in Neonatal Care*, 2, 27-36.
- Rivkees, S.A. (2003). Developing circadian rhythmicity in infants. *Pediatrics*, 112(2), 373-381.
- Rivkees, S.A., et al. (2004). Rest-activity patterns of premature infants are regulated by cycled lighting. *Pediatrics*, 113, 833-839.
- Robinson, J., Moseley, M.J., & Firdler, A.R. (1990). Illuminance of neonatal units. *Archives of Disease and Childhood*, 65(7), 679-682.
- Ross, E.S., & Browne J.V. (2002). Developmental progression of feeding skills: an approach to supporting feeding in preterm infants. *Seminars in Neonatology*, 7, 469-475.
- Shogun, M.G. & Schumann, L.L. (1993). The effect of environmental lighting on the oxygen saturation of preterm infants in the NICU. *Neonatal Network*, 12(5), 7-13.
- Tapia-Rombo, C.A., Morales-Mora, M., & Alvarez-Vazquez, E. (2003). Variations of vital signs, skin color, behavior and oxygen saturation in premature neonates after sponge bathing. Possible complications. *Revista de Investigación Clínica*, 55(4), 438-43.
- Tupekar, R.A., Pinnagoda, J., & Nater, J.P. (1990). The transient and cumulative effect of sodium lauryl sulphate on the epidermal barrier assessed by transepidermal water loss: inter-individual variation. *Acta Dermato-Venerologica*, 70(1), 1-5.
- Vernon H. J., Lane A. T., Wischerath L. J., Davis J. M., & Menegus M. A. (1990). Semipermeable dressing and transepidermal water loss in premature infants. *Pediatrics*, 86(3), 357-362.
- Waber, D., & McCormick, M. (1995). Late neuropsychological outcomes in preterm infants of normal IQ: selective vulnerability of the visual system. *Journal of Pediatric Psychology*, 20, 721-735.
- Wolke, D. (1998). Psychological development of prematurely born children. *Archives of Disease in Childhood*, 78, 567-570.
- Yosipovitch, G., Maayan-Metzger, A., Merlob, P., & Sirota, L. (2000). Skin barrier properties in different body areas in neonates. *Pediatrics*, 106(1 Pt 1), 105-108.



* 1 1906 1032646 09 *

1032646 Develop Progression of Preterm Infants EN Lam (English)
ISBN - 1-59971-437-X